

Salmonella Serotypes - Non-human Isolates 2003

Serotype	Animal													Poultry				Total			
	Avian	Bovine	Canine	Caprine	Cervine	Equine	Feline	Ovine	Porcine	Reptile	Sealion	Meat/bone meal	Environmental	Food*	Spice	Not specified	Neckflap		Feed	Environmental	Miscellaneous including product
Adelaide											7										7
Agona																	1	5	19	3	28
Amsterdam														1							1
Anatum		1				1						13				1			3		19
Anatum 15+															1				1		2
Bovismorbificans		1																			1
Braenderup															1						1
Brandenburg		30	1			6		95				11	11	12	4		8	7	1	186	
Brunei														1							1
Cerro											2										2
Cubana																10					10
Derby											2	2					11		1		16
Enteritidis phage type 9a	2	2	2					1							1		1		1		10
Falkensee 15+													1								1
Give 15+																	3				3
Hadar																1					1
Havana											2	1			1		2	2			8
Heidelberg		3																			3
Hindmarsh		4						66					2		3						75
Infantis	2	1				1		1			2	1	5		34	7	6	4	13		77
Kentucky			1														1				2
Kiambu		1																			1
Lille											9	1									10
London 15+																	7				7
Mbandaka															2		11	4	2		19
Mississippi								1													1
Montevideo											1	3	21		4						29
Newport										6			1								7
Oranienburg		2													1		5				8
Orion					1																1
Orion 15+													3				2				5
Putten																	1				1
Rissen																	4				4
Rubislaw															1		1		1		3
Ruiru																	1				1
Saintpaul	1	5	1	2		1							1								11
Senftenberg			1														3	13			17
Tennessee						2					1	1			1		5	5			15
Thompson																	1	1			2
Typhimurium phage type 1	1	26		1		1						1	1		1		2	26	31		90
8		6						1							1						8
8 variant		2																			2
9		10				1		1				1	1		1		3				18
12a		5										1	2				3	4	7		22
23		5						1					1		1						8
32																	1				1
41																8				1	9
41 variant												1									1
42		8				3		1				32	1		1			1	4		51
42a		1																			1
(N) 74		9		1									1				1				12
101	1	17	1			2		2				2	1		2	3		2	2		35
135	1	18				1		1						10	1	1	6	2			41
135 variant																1					2
155																					1
156		25	3	1					1			7		2		9	3	1			52
160	19	4	1			10	8		2			6	3			11	6	5			75
193		1							1								1				3
199															1						1
RDNC	1	7	2					2	1				1		2						16
Rough		1				1										1	2				5
Untypable		6															1	2	2		11
Weltevreden														2	1	1					4
Westhampton																	1				1
Group B 4,12 :- :-																				1	1
Group B 4,12 :- : 1,2																			1		1
Group B 4,5,12 :- :- (non motile)															1			3			4
Group C 6,7 :- :- (non motile)																1	10		3		14
Group C 6,7 : k :-																		1	1		2
Group C 8,20 : y :-													1								1
Group E 3,15 : e,h :-																	1				1
Group E 3,19 :- :- (non motile)																		2			2
Group F 11 : r :-															1						1
Group I 16 : l,v :-																	1				1
Group R 40 :- :-										1											1
Rough : g,s,t :-															1						1
Rough : r : 1,5								1													1
Rough : s,t :-												1									1
Rough z10 : e,n,z15																	1				1
<b>TOTAL</b>	<b>27</b>	<b>201</b>	<b>13</b>	<b>5</b>	<b>1</b>	<b>28</b>	<b>12</b>	<b>173</b>	<b>5</b>	<b>1</b>	<b>10</b>	<b>48</b>	<b>72</b>	<b>61</b>	<b>1</b>	<b>100</b>	<b>16</b>	<b>126</b>	<b>116</b>	<b>83</b>	<b>1099</b>

S. Typhimurium 465  
 Other serotypes 638  
 Poultry isolates 341  
 Animal isolates 477  
 \* casings and animal carcasses from meatworks are included

(N) S. Typhimurium isolates previously reported as phage type  
 RDNC Aug 01 have now been identified as phage type 74.